

Preparation of manuscripts for the American Journal of Physics using L^AT_EX

Harvey Gould*

Clark University, Department of Physics, Worcester, MA 01610[†]

Jan Tobochnik

Kalamazoo College, Department of Physics, Kalamazoo, MI 49007

(Dated: September 30, 2005)

Abstract

We gave some examples of the use of L^AT_EX that we hope will be helpful in preparing manuscripts for AJP.

I. INTRODUCTION

\TeX looks more difficult than it is. It is almost as easy as π . See how easy it is to make special symbols such as α , β , γ , δ , $\sin x$, \hbar , λ , \dots . We also can make subscripts A_x , A_{xy} and superscripts, e^x , e^{x^2} , and e^{a^b} . We will use \LaTeX , which is based on \TeX and has many higher-level commands (macros) for formatting, making tables, etc. More information can be found in Ref. 1, a book written by Kopka and Daly.¹

We just made a new paragraph. Extra lines and spaces make no difference. Note that all formulae are enclosed by $\$$ and occur in *math mode*.

The default font is Computer Modern. It includes *italics* or *italics*, **boldface** or **boldface**, *slanted* or *slanted*, and `monospaced` or `monospaced` (typewriter) fonts.

II. EQUATIONS

Let us see how easy it is to write equations.

$$\Delta = \sum_{i=1}^N w_i (x_i - \bar{x})^2. \quad (1)$$

It is usually a good idea to number equations, but we can have a equation without a number by writing

$$P(x) = \frac{x - a}{b - a},$$

or

$$g = \frac{1}{2} \sqrt{2\pi}.$$

We can give an equation a label so that we can refer to it later.

$$E = -J \sum_{i=1}^N s_i s_{i+1}, \quad (2)$$

Equation (2) expresses the energy of a configuration of spins.² If we use the `amsmath` package, we could write Eq. (2).

We can define our own macros to save typing. For example, suppose that we introduce the macros:

```
\newcommand{\lb}{\langle}
```

```
\newcommand{\rb}{\rangle}
```

Then we can write the average value of x as

```
\begin{equation}
\lb x \rb = 3
\end{equation}
```

The result is

$$\langle x \rangle = 3. \tag{3}$$

Examples of more complicated equations:

$$I = \int_{-\infty}^{\infty} f(x) dx. \tag{4}$$

We can do some fine tuning by adding small amounts of horizontal spacing:

```
\, small space      \! negative space
```

as is done in Eq. (4).

We also can align several equations:

$$a = b \tag{5}$$

$$c = d, \tag{6}$$

or number them as subequations:

$$a = b \tag{7a}$$

$$c = d. \tag{7b}$$

How would you suppress the equation numbers in these two examples? A slightly better way to align equations use the `amsmath` package:

$$a = b \tag{8a}$$

$$c = d. \tag{8b}$$

Can you notice the difference? Some other examples:

$$m(T) = \begin{cases} 0 & T > T_c \\ (1 - [\sinh 2\beta J]^{-4})^{1/8} & T < T_c \end{cases} \tag{9}$$

$$\begin{aligned} \mathbf{T} &= \begin{pmatrix} T_{++} & T_{+-} \\ T_{-+} & T_{--} \end{pmatrix} \\ &= \begin{pmatrix} e^{\beta(J+B)} & e^{-\beta J} \\ e^{-\beta J} & e^{\beta(J-B)} \end{pmatrix} \end{aligned} \tag{10}$$

$$\sum_i \vec{A} \cdot \vec{B} = -P \int \mathbf{r} \cdot \hat{\mathbf{n}} dA = P \int \vec{\nabla} \cdot \mathbf{r} dV. \tag{11}$$

III. LISTS

Some example of formatted lists include the following:

1. bread
2. cheese
 - Tom
 - Dick

An example of a table is given in Table I at the end of the manuscript and examples of how to include figures are shown in Figs. 1 and 2.

IV. SPECIAL SYMBOLS

A. Common Greek letters

These commands may be used only in math mode. Only the most common letters are included.

$\alpha, \beta, \gamma, \Gamma, \delta, \Delta, \epsilon, \zeta, \eta, \theta, \Theta, \kappa, \lambda, \Lambda, \mu, \nu, \xi, \Xi, \pi, \Pi, \rho, \sigma, \tau, \phi, \Phi, \chi, \psi, \Psi, \omega, \Omega$

B. Special symbols

The derivative is defined as

$$\frac{dy}{dx} = \lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x} \tag{12}$$

$$f(x) \rightarrow y \quad \text{as } x \rightarrow x_0 \quad (13)$$

$$f(x) \xrightarrow{x \rightarrow x_0} y \quad (14)$$

Order of magnitude:

$$\log_{10} f \simeq n \quad (15)$$

$$f(x) \sim 10^n \quad (16)$$

Approximate equality:

$$f(x) \simeq g(x) \quad (17)$$

TeX is simple if we keep everything in proportion:

$$f(x) \propto x^3. \quad (18)$$

We can skip some space by using a command such as

`\bigskip` `\medskip` `\smallskip` `\vspace{1pc}`

The space can be negative.

And it sometimes is convenient to write \tilde{x} , \widetilde{xy} , \overline{A} , $\%$, accents: Schrödinger and é (or any other letter \acute{z}).

V. LITERAL TEXT

It is desirable to print program code exactly as it is typed in a monospaced font. Use `\begin{verbatim}` and `\end{verbatim}` as in the following example:

```
public void computeArea()
{
    this.area = this.length*this.length;
    System.out.println("Area = " + this.area);
}
```

The command `\verbatiminput{programs/Square.java}` allows the file `Square.java` in the direction `programs` to be listed without changes.

APPENDIX A: SOME DOS AND DON'TS

1. Note the AJP style for books¹ and articles.³
2. Also note the American convention of the positions of citations.
3. Do not skip a line before `\begin{equation}` or after `\end{equation}`.

ACKNOWLEDGMENTS

Be sure to thank your colleagues and any granting agencies.

* Also at home.

† Electronic address: hgould@clarku.edu

¹ Helmut Kopka and Patrick W. Daly, *A Guide to L^AT_EX: Document Preparation for Beginners and Advanced Users* (Addison-Wesley, Reading, MA, 1999), 3rd. ed.

² It is necessary to process a file twice to get the counters correct. AJP does not use footnotes.

³ B. C. Freasier, C. E. Woodward, and R. J. Bearman, "Heat capacity extrema on isotherms in one-dimension: Two particles interacting with the truncated Lennard-Jones potential in the canonical ensemble," *J. Chem. Phys.* **105**, 3686–3690 (1996).

TABLES

Tables are a little difficult until you get the knack. \LaTeX automatically calculates the width of the columns. Tables should be placed at the end of the manuscript.

lattice	d	q	T_{mf}/T_c
square	2	4	1.763
triangular	2	6	1.648
diamond	3	4	1.479
simple cubic	3	6	1.330
bcc	3	8	1.260
fcc	3	12	1.225

TABLE I: Comparison of the mean-field predictions for the critical temperature of the Ising model with exact results and the best known estimates for different spatial dimensions d and lattice symmetries.

FIGURES

It is easy to include encapsulated postscript files (see Figure 1). We can make figures bigger or smaller by scaling them. Figure 2 has been scaled by 80%. Figures should be placed at the end of the manuscript and sent as separate files. It also is possible to include pdf files.




FIG. 1: Show me a sine.




FIG. 2: Plot of the Lennard-Jones potential $u(r)$. The potential is characterized by a length σ and an energy ϵ . This potential is applied in Ref. 3.